

Intergenic and External Transcribed Spacers of Ribosomal RNA Genes in Lizard-infecting *Leishmania*: Molecular Structure and Phylogenetic Relationship to Mammal-infecting *Leishmania* in the Subgenus *Leishmania* (*Leishmania*)

Tereza C Orlando, Mary Anne T Rubio*, Nancy R Sturm*, David A Campbell*,
Lucile M Floeter-Winter/⁺

Departamento de Parasitologia, Instituto de Ciências Biomédicas, Universidade de São Paulo, Av. Prof. Lineu Prestes 1374, 05508-900 São Paulo, SP, Brasil *Department of Microbiology, Immunology & Molecular Genetics, University of California, Los Angeles, USA

To establish the relationships of the lizard- and mammal-infecting *Leishmania*, we characterized the intergenic spacer region of ribosomal RNA genes from *L. tarentolae* and *L. hoogstraali*. The organization of these regions is similar to those of other eukaryotes. The intergenic spacer region was approximately 4 kb in *L. tarentolae* and 5.5 kb in *L. hoogstraali*. The size difference was due to a greater number of 63-bp repetitive elements in the latter species. This region also contained another element, repeated twice, that had an inverted octanucleotide with the potential to form a stem-loop structure that could be involved in transcription termination or processing events. The ribosomal RNA gene localization showed a distinct pattern with one chromosomal band (2.2 Mb) for *L. tarentolae* and two (1.5 and 1.3 Mb) for *L. hoogstraali*. The study also showed sequence differences in the external transcribed region that could be used to distinguish lizard *Leishmania* from the mammalian *Leishmania*. The intergenic spacer region structure features found among *Leishmania* species indicated that lizard and mammalian *Leishmania* are closely related and support the inclusion of lizard-infecting species into the subgenus *Sauroleishmania* proposed by Saf'janova in 1982.

Key words: lizard leishmania - RNA polymerase I - rRNA processing - repetitive DNA - rRNA spacer evolution

Leishmania tarentolae and *L. hoogstraali* are blood-stream kinetoplastids of lizards. Their geographical distribution is restricted to the Old World. Lizard-infecting *Leishmania* are closely related to mammalian-infecting *Leishmania*, but their exact taxonomic position and phylogenetic relationship remains uncertain. Recently, biochemical and biological data have been used in comparative studies of lizard and mammalian *Leishmania* species (Simpson & Holz 1988, Briones et al. 1992, Fu & Kolesnikov 1994, Croan & Ellis 1996, Previato et al. 1997, Noyes et al. 1997, Croan et al. 1997, Shaw 1997, Noyes et al. 1998) indicating a close relationship.

Lizard-infecting *Leishmania* play an important role in epidemiological studies of leishmaniasis. Amastigote forms of *L. adleri* (another lizard *Leishmania* species) were detected in volunteers at the site of inoculation five

days after infection with *L. adleri* promastigotes. Serological studies also showed common antigens among *L. adleri* and *L. hoogstraali* and some mammalian *Leishmania* species. This observation has an implication in the kala-azar epidemiology since false positives could be generated due to naturally-transmitted *L. adleri* (see revision in Wilson & Southgate 1979, Telford 1995). In addition, amastigote forms have been detected in phagocytic cells of naturally and experimentally infected lizards (Dollahon & Janovy 1974, Elwasila 1988).

Earlier studies using the small subunit ribosomal RNA (rRNA) molecule helped to establish the lizard *Leishmania* proximity to mammalian *Leishmania* (Briones et al. 1992) relative to other trypanosomatid genera. However, the small subunit sequences are not suitable for resolution of species within the same genera. On the other hand, the rRNA gene intergenic spacer region (IGS) and the external transcribed spacer (ETS) are under less selective pressure and thus provide a means to distinguish evolutionary relationships among more closely-related species. The IGS/ETS region of trypanosomatids have been widely studied (White et al. 1986, Grondal et al. 1990, Zomerdijk et al. 1991, Rudenko et al. 1991, Dietrich et al. 1993, Martinez-Calvillo & Hernandez 1994, Tyler-Cross et al. 1995, Uliana et al. 1996, Gay et al. 1996, Yan et al. 1999, Downey & Donelson 1999, Schnare et al. 2000) and shows a general organization similar to the majority of eukaryotes.

In the present work, the molecular structure of the IGS region from two lizard-infecting *Leishmania* species is described and compared to the corresponding sequences

This work was supported by Fapesp (Fundação de Amparo à Pesquisa do Estado de São Paulo), CNPq and NIH (Grant no. AI34536). TCO was supported by a fellowship from Fapesp and MATR was supported by a National Science Foundation Graduate Research Fellowship.

Note: Nucleotide sequence data reported here are available in the GenBank database under the accession numbers AY026525 and AF345631-33.

*Corresponding author: Fax: +55-11-3818.7329. E-mail: lmfwinte@icb.usp.br

Received 13 September 2001

Accepted 6 February 2002

from the mammal-infecting *Leishmania*. These data refine the phylogenetic positioning of the lizard *Leishmania* and support their status as a subgenus of *Leishmania*.

MATERIALS AND METHODS

Organisms - Promastigotes of *L. tarentolae* (ATCC 30267) and *L. hoogstraali* (RHEM/SD/1963/NG26) from the Trypanosomatid Bank at Universidade de São Paulo were cultivated at 25°C in LIT medium (Camargo 1964).

DNA and RNA extractions, isolation of the IGS/ETS region and sequencing - Genomic DNA from *Leishmania* species was purified as previously described (Uliana et al. 1991). Total RNA was extracted as described (Chomczynski & Sacchi 1987).

A genomic *ApaI*-*BglII* fragment encompassing ~3 kb of the IGS/ETS region of *L. tarentolae* was isolated from a cosmid library (Fig. 1C, clone pAB 2.8). Clone pL1.3 (Fig. 1C) was obtained by the isolation of an *AluI* fragment from pAB 2.8. For *L. hoogstraali*, a 9-kb *BglIII* fragment was isolated from a partial genomic library, constructed using inserts of 7 to 10 kb produced by a complete digestion of genomic DNA with *BglII*. This fragment contains ~2 kb from the large subunit rRNAs subunits and the whole IGS/ETS region. Fig. 1B shows clones pLh 6500, pLh 3400 and pLh 8A, containing small fragments, used in this work. These clones were mapped with several restriction enzymes, subcloned into pUC 18/19 series (Fig. 1) and sequenced by the dideoxynucleotide method (Sanger et al. 1977). The sequences were determined by a combination of manual and automated sequencing using an ABI 377 or 310 sequencer (Perkin Elmer).

Chromosomal analysis - Chromosome bands were separated by pulsed-field gel electrophoresis (Schwartz & Cantor 1984) in 1.2% agarose gels and 0.5x TBE buffer. The running conditions were: three phases pulses of 180, 240 and 300 s at 6 V/cm for 18 h each pulse at 10°C. Blots were probed with the *AccI* fragment (Fig. 1C) and with the small subunit rRNA from *L. tarentolae*.

Sequence and phylogenetic analysis - The sequences were aligned with CLUSTALW 1.6 (Thompson et al. 1994) and the matrix obtained contained 1119 characters and 6 taxa. *Crithidia fasciculata* was used as the outgroup.

Phylogenetic trees were constructed using PAUP version 4.03b (Swofford 1999) for distance and parsimony methods. Maximum likelihood trees were obtained with BASEML program from the PAML 2.0 package (Yang 1999) using the K80 model of DNA substitution with gamma distribution ($\alpha = 0.85$). Bootstrap values were calculated for 100 replicates.

RESULTS

IGS/ETS sequence comparison - Based in a Northern blot experiment the approximate size of the ETS region of *L. hoogstraali* and *L. tarentolae* was determined (data not shown) and used to position the data to align and compare this region with the corresponding sequences of *L. (L.) amazonensis*, *L. (L.) chagasi*, *L. (L.) donovani* and *C. fasciculata* (Fig. 2, Table I). The comparison showed blocks of sequence similarity among the *Leishmania* species and *Crithidia*. Motifs were identified corresponding to those described previously in *L. (L.) amazonensis* (Uliana et al. 1996) and two regions (Fig. 2) of absolute identity to the proposed base-pairing sites of




Fig. 1: scheme of the general ribosomal RNA gene arrangement (A) with the respective intergenic spacer/external transcribed spacer region of *Leishmania hoogstraali* (B) and *L. tarentolae* (C). The blank box in B represents the 3' end of the rRNA gene 24Sβ. A, *ApaI*; B, *BamHI*; C, *HincII*; D, *HindIII*; G, *BglIII*; J, *AccI*; P, *PstI*; S, *SmaI*; U, *AluI*; V, *PvuII*; X, *XhoI*; Y, *DdeI*. The 63 bp repetitive elements are represented in stippled and white rectangles, the flag maps the estimated transcription start point, the black rectangles indicate the element containing the inverted octanucleotides. The thick line corresponds to the sequenced region.

the U3 small nucleolar RNA (snoRNA) of *C. fasciculata* (Schnare et al. 2000) and *L. tarentolae* (Shi et al. 1994). A *Hind*III site in the estimated ETS region of *L. tarentolae* and *L. hoogstraali* (Fig. 1) could be used to discriminate lizard isolates from the mammalian isolates.

L. tarentolae and *L. hoogstraali* IGS/ETS structure - The IGS/ETS structure was determined by sequence analysis of the corresponding genomic segments from each species (Fig. 1): ~4 kb for *L. tarentolae* and 5.5 kb for *L. hoogstraali*.

The relevant IGS/ETS fragments were digested with restriction enzymes that had 4-bp recognition sites. The patterns obtained for both species with *Dde*I revealed the presence of repetitive element in both regions. A non-stoichiometric quantity of a ~70-bp fragment accumulated with partial-restriction kinetics, and the sum of all frag-

ments generated by complete digestion was less than the size of the original fragment (data not shown).

Determining the nucleotide sequence of the IGS fragments allowed us to fully characterize the repetitive elements. For both species, the first 63-bp repeat is located ~200 bp upstream of the estimated transcription start point (Fig. 1B, C, hatched rectangles). By sequence analysis we detected approximately 12 repeats for *L. tarentolae* and 40 for *L. hoogstraali*.

These repetitive elements showed approximately 90% sequence identity between lizard *Leishmania* species (Table I, upper section). In contrast, comparison with the same repetitive elements in the IGS of mammalian *Leishmania* showed a very low degree of similarity (Table I, upper section). Fig. 3 shows a sequence comparison of the repetitive elements found in each species.




Fig. 2: sequence comparison of a segment of the external transcribed spacer. AMA corresponds to *Leishmania* (*L.*) *amazonensis*, CHA corresponds to *L.* (*L.*) *chagasi*, DON corresponds to *L.* (*L.*) *donovani*, TAR corresponds to *L. tarentolae*, HOO corresponds to *L. hoogstraali* and CRIT corresponds to *Crithidia fasciculata*. GeneBank accession numbers for *L.* (*L.*) *amazonensis*, *L.* (*L.*) *chagasi*, *L.* (*L.*) *donovani* and *C. fasciculata* are U21687, U42465, L38572 and Y00055, respectively. Regions I and II correspond to the proposed site of base pairing with the U3 snoRNA in *C. fasciculata* (Schnare et al. 2000) and *L. tarentolae* (Shi et al. 1994). Numbering is in relation to the transcription start point of *L. amazonensis* (Uliana et al. 1996). Dots (.) represent nucleotide identity, dashes (-) represent gaps introduced to improve the alignment obtained with Clustalw (1.60) program.

TABLE

Percentage of similarity among the six external transcribed spacer sequences (lower part) and in the 63bp repetitive elements (upper part)

	<i>L. tarentolae</i>	<i>L. hoogstraali</i>	<i>L.</i> (<i>L.</i>) <i>amazonensis</i>	<i>L.</i> (<i>L.</i>) <i>chagasi</i>	<i>L.</i> (<i>L.</i>) <i>donovani</i>
<i>Leishmania tarentolae</i>	-	90	36	34	36
<i>L. hoogstraali</i>	93	-	35	34	36
<i>L.</i> (<i>L.</i>) <i>amazonensis</i>	85	87	-	68	68
<i>L.</i> (<i>L.</i>) <i>chagasi</i>	85	86	88	-	96
<i>L.</i> (<i>L.</i>) <i>donovani</i>	86	88	90	97	-
<i>Crithidia fasciculata</i>	60	62	59	58	59

```

HOO  TGTGCAGCTACGCGGGCACGCTCAGCGAGGCATTGTCTGGCGTCGCAGCAGTGCAGTATTGGC-
TAR  T . . G . AGCCATGCAGG . ACGC . . AGTGA . GCATT . T . TGG . . CNGCAGT . G . G . GGTATTGGC-
AMA  A . . A . TTGGACATTCC . --TT . . GCCAC . CCAGA . - . CGT . . -GAGGCC . C . C . TTCGCGTCCC
CHA  A . . A . TGGAGCATTCC . GTTT . . GCTGC . AGGAA . A . CCT . . CGGGAGC . T . G . TTCGCGCCCC
DON  A . . A . TGGAGCATTCCG . GTTT . . GCTGC . AGGGA . A . CCT . . CGGGAGC . T . G . TTCGCGCCCC

```

Fig. 3: repetitive elements found in the intergenic spacer region of the rRNA genes of *Leishmania* species. Species identification as in Fig. 2. Bold type is the *DdeI* site. Dots (.) represent nucleotide identity, dashes (-) represent gaps introduced to improve the alignment obtained with Clustalw (1.60) program.

The results of the probed pulsed-field gel electrophoresis blots (Fig. 4) indicated that the repetitive elements co-localized to the same chromosomal bands as the rRNA of both species and were not dispersed throughout the genome. The pulse-field pattern also revealed that the rRNA gene maps to chromosomal bands of different size (Fig. 4) in *L. tarentolae* (2.2 Mb) and *L. hoogstraali* (1.5 and 1.3 Mb). Hybridization experiments using high stringency conditions indicated the presence of these elements in *L. adleri* and *L. gymnodactyli* (other lizard *Leishmania* species). The 63-bp repetitive elements from lizard *Leishmania* did not hybridize with *L. (L.) amazonensis* nor *L. (L.) mexicana* chromosomal bands (data not shown).

Upstream from the 63-bp repetitive elements, an inverted octanucleotide motif repeated twice was found in both species (Fig. 1B, C, black rectangles). A similar sequence was described in the same IGS position in *L. (L.) infantum* (Requeña et al. 1997) and recently, it was also found in the IGS region of *C. fasciculata* (Schnare et al. 2000). A comparison among these sequences showed that the repeats in *C. fasciculata* and *Leishmania* species are not as conserved as among the *Leishmania* species repeats (Fig. 5A). The potential to form a stem-loop struc-

ture as identified in the *L. (L.) infantum* sequence was also found for the two blocks of *L. tarentolae*, *L. hoogstraali* and *C. fasciculata* (Fig. 5B).

Phylogeny - Trees derived from distance, parsimony and maximum likelihood (Fig. 6) methods exhibited identical topologies and showed that the *Leishmania* of lizards studied are very close to each other. The lizard *Leishmania* form a sister group to the mammalian *Leishmania* of the subgenus *L. (Leishmania)* analyzed with *C. fasciculata* used as the outgroup. The bootstrap values obtained for each method (Fig. 6, nodes) support the hypothesis that lizard *Leishmania* and mammalian *Leishmania* species of the subgenus *L. (Leishmania)* are sister groups.

DISCUSSION

In this study we have determined the sequence of the IGS/ETS upstream from the small subunit rRNA of two lizard-infecting *Leishmania* species for comparison with their mammal-infecting relatives. The size of the IGS region determined for both species is approximately the same as the size found for *L. (L.) infantum* (5 kb, Requeña et al. 1997) and *L. (L.) donovani* (4 kb, Lodes et al. 1995), indicating that the size of this region is conserved between




Fig. 4: chromosomal pattern of *Leishmania tarentolae* (1 and 3) and *L. hoogstraali* (2 and 4) rRNA gene localization. A: chromosomal bands after ethidium bromide staining. M corresponds to Megabase IV (Invitrogen); B: autoradiogram of the chromosome blot in panel A. In panel I the fragment *AccI/AccI* from *L. tarentolae* (Fig. 1A) was used as the probe and in panel II the 18S subunit DNA was used as the probe. Washing conditions: 2x SSC/0.1% SDS at 45°C. Exposure time: 5 days at -20°C for panel I and 4 days at -20 °C for panel II.




Fig. 5: analysis of the three repetitive elements found downstream from the *Leishmania* (*L.*) *infantum* large subunit rRNA (INF 1, 2 and 3), *L. tarentolae* (TAR 4 and 5), *L. hoogstraali* (HOO 6 and 7) and *Crithidia fasciculata* (CRIT). A: alignment of the elements. Bold letters correspond to the inverted octanucleotides. Asterisks (*) below the alignment indicate base identity in each position among the eight sequences; dashes (-) represent gaps introduced to maximize the alignment; B: sequences shown in A can form similar stem loop structures. I *L. (L.) infantum* (Requeña et al. 1997); II. *L. tarentolae*. III. *L. hoogstraali*. IV. *C. fasciculata* (from R3 sequence described in Schnare et al. 2000).




Fig. 6: topology obtained with the maximum likelihood method for the external transcribed spacer sequence alignment. Distance method (performed with the neighbor-joining algorithm) and parsimony method gave similar topologies, although the bootstrap values obtained differed: numbers at the left of each node represent maximum likelihood values, numbers at the right represent neighbor-joining values and numbers under each node represent parsimony values. The bar indicates distance for maximum likelihood.

lizard and mammalian *Leishmania* species; the size observed in other eukaryotes is variable. The differences in length can be attributed to a variable number of 63-bp repetitive-element found in each lizard species.

The estimated ETS region of each species (~1000 bp) was shown to be very similar in size to *L. (L.) amazonensis* (1050 bp, Uliana et al. 1996), *L. (L.) chagasi* (1060 bp, Gay et al. 1996), *L. (L.) donovani* (1020 bp, Yan et al. 1999), *C. fasciculata* (1050 bp, Grondal et al. 1990) and *T. congolense* (961 bp, Downey & Donelson 1999). For *T. cruzi* and *T. brucei* the ETS size is significantly larger [1550 bp, Dietrich et al. (1993) and 1200 bp, White et al. (1986), respectively].

The comparison of ETS regions showed conserved motifs between lizard and mammalian *Leishmania* species and *C. fasciculata* (data not shown), including the proposed sites of base-pairing with the U3 snoRNA of *C. fasciculata* (Schnare et al. 2000) and *L. tarentolae* (Shi et al. 1994) (Fig. 2). The presence of similar motifs suggests a common processing mechanism of the pre-rRNA molecule.

As in many other eukaryotes, conserved, repeated elements upstream of the predicted transcription start point were detected in both species. However, when compared to mammalian *Leishmania* species, the degree of similarity dropped three fold (Table I). The repeat number varied among the *Leishmania* species: two populations of 35 and 70 copies in *L. (L.) amazonensis* (Uliana et al. 1996)

and 39 copies in *L. (L.) donovani* (Yan et al. 1999), while for lizard *Leishmania* approximately 12 copies of the 63-bp repeat in the *L. tarentolae* IGS and approximately 40 copies in *L. hoogstraali* were found. The size of the repeated elements was conserved in *Leishmania* (~ 60 bp), while 28 copies of a 19-bp element were found located upstream of the transcription start point in *C. fasciculata* (Schnare et al. 2000), indicating that the repeat size is not necessarily conserved in the kinetoplastids. In *L. (L.) chagasi* these elements showed a modest enhancer activity in a transient transfection assay (Gay et al. 1996).

The chromosomal analysis (Fig. 4) showed that the rRNA genes and the 63-bp repetitive elements mapped to the same chromosomal bands for both species. However, the detection of two chromosomal bands of different size in *L. hoogstraali* (1.3 and 1.5 Mb) does not discriminate whether they reside on homologous or different chromosomes. A similar pattern of two chromosomal bands for the rRNA genes was also identified for some *Viannia* subgenus species (Inga et al. 1998), as well as in *L. (L.) donovani* (Yan et al. 1999).

The analysis of an element located downstream of the large subunit 2 (24S β) of *L. tarentolae*, *L. hoogstraali*, *L. (L.) infantum* (Requeña et al. 1997) and *C. fasciculata* (Schnare et al. 2000) showed high degrees of similarity, mainly in the inverted octanucleotides (Fig. 5A, bold) and in the nucleotides that can form potential stem-loop structures (Fig. 5B).

This conservation indicates a possible biological role to this element, probably in the transcription termination of pre-rRNA or in subsequent processing events. The sequence of the element is conserved among *Leishmania* species, demonstrating the close relationship between lizard and mammalian *Leishmania*. An intriguing aspect of these motifs is that they have inverted octanucleotides identical to blocks found in λ phage and *Escherichia coli* (Thaler & Stahl 1988), where they define hot spots for recombination. Thus, the repeated sequences could be involved in recombination events, as evidenced by the correlation of the repetitive sequences to the heterogeneity in the copy number of the rRNA genes.

The alignment of ETS sequences showed a high degree of sequence similarity between lizard *Leishmania* and mammal-infecting *Leishmania* (Table, lower section). The analysis of the data using diverse phylogenetic approaches resulted in the generation of identical trees showing that the lizard *Leishmania* are very closely related to *L. (L.) donovani*. These results are in agreement with biogeographic data, since both lizard *Leishmania* and *L. (L.) donovani* are found only in the Old World (Shaw 1994). Similar phylogenetic relationships for these organisms were found using the DNA polymerase I genes and RNA polymerase II genes of different *Leishmania* species (Croan et al. 1997). Furthermore, functional results involving the exact transcription starting point, the ETS and the promoter region of the rRNA gene of lizard *Leishmania* showed some degree of heterologous recognition of these regions between lizards and mammalian-infecting species (Orlando et al., manuscript in preparation), corroborating their close relationship also in a functional level, as this recognition is thought to be highly species-specific (Grummt et al. 1982, Sommerville 1984).

Both the molecular data presented and the inferred phylogenetic relationships are in agreement with the inclusion of the lizard *Leishmania* in the subgenus *Sauroleishmania*, as proposed by Saf'janova, using biological criteria (Saf'janova 1982).

ACKNOWLEDGEMENTS

To Jeffrey J Shaw for critical reading and comments on the manuscript and Sergio Pereira for help with the phylogeny.

REFERENCES

- Briones MRS, Nelson K, Beverley SM, Affonso MHT, Camargo EP, Floeter-Winter LM 1992. *Leishmania tarentolae*: taxonomic relatedness inferred from phylogenetic analysis of small subunit ribosomal RNA gene. *Mol Biochem Parasitol* 53: 121-128.
- Camargo EP 1964. Growth and differentiation of *Trypanosoma cruzi*. I. Origin of metacyclic trypanosomes in liquid media. *Rev Inst Med Trop São Paulo* 6: 93-100.
- Chomczynski P, Sacchi N 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162: 156-159.
- Croan DA, Ellis J 1996. Phylogenetic relationship between *Leishmania*, *Viannia* and *Sauroleishmania* inferred from comparison of a variable domain within the RNA polymerase II largest subunit gene. *Mol Biochem Parasitol* 79: 97-102.
- Croan DA, Morrison DA, Ellis J 1997. Evolution of the genus *Leishmania* revealed by comparison of DNA and RNA polymerase gene sequences. *Mol Biochem Parasitol* 89: 149-159.
- Dietrich P, Soares MBM, Affonso MHT, Floeter-Winter LM 1993. *Trypanosoma cruzi* ribosomal gene: analysis of promoter and upstream intergenic spacer sequences. *Gene* 125: 103-107.
- Dollahon NR, Janovy Jr J 1974. Experimental infection of New World lizards with Old World lizard *Leishmania* species. *Exp Parasitol* 36: 253-260.
- Downey N, Donelson JE 1999. Search for promoters for GARP and rRNA genes of *Trypanosoma congolense*. *Mol Biochem Parasitol* 104: 25-38.
- Elwasila M 1988. *Leishmania tarentolae* Wenyon, 1921 from the gecko *Tarentola annularis* in the Sudan. *Parasitol Res* 74: 591-592.
- Fu O, Kolesnikov AA 1994. *Leishmania gymnodactyli* and *Leishmania infantum* minicircles contain the same guide RNA genes as do minicircles of *Leishmania tarentolae*. *Mol Biochem Parasitol* 67: 171-174.
- Gay LS, Wilson ME, Donelson JE 1996. The promoter for ribosomal RNA genes of *Leishmania chagasi*. *Mol Biochem Parasitol* 77: 115-252.
- Grondal EJM, Evers R, Cornelissen AWCA 1990. Identification and sequence analysis of the ribosomal DNA promoter region of *Crithidia fasciculata*. *Nucleic Acids Res* 18: 1333-1338.
- Grummt I, Roth E, Paule MR 1982. Ribosomal RNA transcription in vitro is species specific. *Nature* 296: 173-174.
- Inga R, De Doncker S, Gomez J, Lopez M, Garcia R, Le Ray D, Arevalo J, Dujardin JC 1998. Relation between variation in copy number of ribosomal RNA encoding genes and size of harbouring chromosomes in *Leishmania* of subgenus *Viannia*. *Mol Biochem Parasitol* 92: 219-228.
- Lodes MJ, Merlin G, Devos T, Ghosh A, Madhubala R, Myler PJ, Stuart K 1995. Increased expression of LD1 genes transcribed by RNA polymerase I in *Leishmania donovani* as a result of duplication into the rRNA gene locus. *Mol Cell*

- Biol* 15: 6845-6853.
- Martinez-Calvillo S, Hernandez R 1994. *Trypanosoma cruzi* ribosomal DNA: mapping of a putative distal promoter. *Gene* 142: 243-247.
- Noyes HA, Arana BA, Chance ML, Maingon R 1997. The *Leishmania hertigi* Kinetoplastida; Trypanosomatidae complex and the lizard *Leishmania*: their classification and evidence for a neotropical origin of the *Leishmania-Endotrypanum* clade. *J Eukaryot Microbiol* 44: 511-517.
- Noyes HA, Chance ML, Croan DG, Ellis JT 1998. *Leishmania (Sauroleishmania)*: a comment on classification. *Parasitol Today* 14: 167.
- Previato JO, Jones C, Wait R, Routier F, Saraiva E, Mendonça-Previato L 1997. *Leishmania adleri*, a lizard parasite, expresses structurally similar glycoinositolphospholipids to mammalian *Leishmania*. *Glycobiol* 7: 687-695.
- Requeña MJ, Soto M, Quijada L, Carrillo G, Alonso C 1997. A region containing repeated elements is associated with transcriptional termination of *Leishmania infantum* ribosomal RNA genes. *Mol Biochem Parasitol* 84: 101-110.
- Rudenko G, Chung H-MM, Pham VP, Van Der Ploeg LHT 1991. RNA polymerase I can mediate expression of CAT and neo protein-coding genes in *Trypanosoma brucei*. *EMBO J* 10: 3387-3397.
- Saf'janova VM 1982. [The problem of taxonomy with *Leishmania*]. *Ser Protozool Sov Acad Sci Leningr* 7: 5-109.
- Sanger F, Nicklen S, Coulson AR 1977. DNA sequencing with chain terminating inhibitors. *Proc Natl Acad Sci USA* 74: 5463-5467.
- Schnare MN, Collings JC, Spencer DF, Gray MW 2000. The 28S-18S rDNA intergenic spacer from *Crithidia fasciculata*: repeated sequences, length heterogeneity, putative processing sites and potential interactions between U3 small nuclear RNA and the ribosomal RNA precursor. *Nucleic Acids Res* 28: 3452-3461.
- Schwartz DC, Cantor CR 1984. Separation of yeast chromosome-sized DNAs by pulsed field gel electrophoresis. *Cell* 37: 67-75.
- Shaw JJ 1997. Ecological and evolutionary pressures on leishmanial parasites. *Braz J Genet* 20: 123-128.
- Shaw JJ 1994. Taxonomy of the genus *Leishmania*: present and future trends and their implications. *Mem Inst Oswaldo Cruz* 89: 471-478.
- Shi X, Chen DT, Suyama Y 1994. A nuclear tRNA gene cluster in the protozoa *Leishmania tarentolae* and differential distribution of nuclear-encoded tRNAs between the cytosol and mitochondria. *Mol Biochem Parasitol* 65: 23-37.
- Simpson L, Holz G 1988. The status of *Leishmania tarentolae/ Trypanosoma platyductyli*. *Parasitol Today* 4: 115-118.
- Sommerville J 1984. RNA polymerase I promoters and transcription factors. *Nature* 310: 189-190.
- Swofford DL 1999. PAUP* Phylogenetic Analysis Using Parsimony* and other methods. Version 4. Sinauer Associates, Sunderland, Massachusetts.
- Telford Jr SR 1995. The Kinetoplastid hemoflagellates of reptiles. In JP Kreier, *Parasitic Protozoa*, Vol. 10, 2nd ed., Academic Press, New York, p. 161-223.
- Thaler DS, Stahl FW 1988. DNA double-chain breaks in recombination of phage and of yeast. *Ann Rev Genet* 22: 169-197.
- Thompson JD, Higgins DG, Gibson TJ 1994. CLUSTALW: improving the sensitivity of progressive multiple sequence alignments through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Res* 22: 4673-4680.
- Tyler-Cross RE, Harris-Kerr C, Floeter-Winter LM, Buck GA 1995. Transient expression mediated by the *Trypanosoma cruzi* rRNA promoter. *Mol Biochem Parasitol* 72: 23-31.
- Uliana SRB, Affonso MHT, Camargo EP, Floeter-Winter LM 1991. *Leishmania*: genus identification based on a specific sequence of the 18S ribosomal RNA sequence. *Exp Parasitol* 72: 157-163.
- Uliana SRB, Fischer W, Stempliuk VA, Floeter-Winter LM 1996. Structural and functional characterization of *Leishmania amazonensis* ribosomal RNA promoter. *Mol Biochem Parasitol* 76: 245-255.
- White TC, Rudenko G, Borst P 1986. Three small RNAs within the 10 kB trypanosome rRNA transcription unit are analogous to Domain VII of other eukaryotic 28S rRNA. *Nucleic Acids Res* 14: 9472-9489.
- Wilson VCLC, Southgate BA 1979. Lizard *Leishmania*. In WHR Lumsden & DA Evans (eds), *Biology of the Kinetoplastida*, Vol. 2, Academic Press, London and New York, p. 241-268.
- Yan S, Lodes MJ, Fox M, Myler PJ, Stuart K 1999. Characterization of the *Leishmania donovani* ribosomal RNA promoter. *Mol Biochem Parasitol* 103: 197-210.
- Yang Z 1999. Phylogenetic analysis by maximum likelihood PAML. Version 2.0. University of College London, program distributed by the author.
- Zomerdiijk JCBM, Kieft R, Shiels PG, Borst P 1991. Alpha-amanitin-resistant transcription units in trypanosomes: a comparison of promoter sequences for a VSG gene expression site and for the ribosomal RNA genes. *Nucleic Acids Res* 19: 5153-5158.